

Case Study: Supplemental Light for Lactating Dairy Cows[†]

M. J. VANBAALE^{1*}, D. V. ARMSTRONG¹, M. A. ETCHEBARNE², R. M.
MATTINGLY², AND J. B. FISCALINI²

¹*Department of Animal Sciences, The University of Arizona, Tucson 85721 and* ²*Fiscalini Dairy Farm, Modesto, CA 95358*

Corresponding Author¹:

Matthew J. VanBaale
The University of Arizona
212 Shantz Building PO Box 210038
Tucson, AZ 85721-0038
Phone: (520) 621-1923
Fax: (520) 621-9435
Email: vanbaale@ag.arizona.edu

This work was partially supported by the University of Arizona Experiment Station #AZRT-137024-R-24-132.

ABSTRACT

Ninety-eight multiparous and 60 primiparous Holsteins were utilized in an extended lighting trial to investigate effects of photoperiod on milk yield. After parturition, all animals were housed in one pen until 20 d in milk (**DIM**) under normal daylight/nightlight conditions. On d 21, animals were randomly assigned to two treatment groups receiving supplemental light (**SL**) or normal daylight and darkness (**NL**). While in freestalls all cows in the SL groups were exposed to 17 h of natural light plus supplemental light above 15-foot candle (**FC**) and 7 h of light below 5 FC in the freestall area. The light exposure for the NL groups followed the normal sunrise-sunset pattern for the north 40th parallel of sunrise 0530 to 0730 and sunset 1700 to 1900, an average of 12 h of light. There was no milk production difference observed between primiparous animals assigned to SL or NL (33.0 vs. 32.9 kg/d; $P = 0.81$) treatments. However, multiparous cows in SL groups produced more milk (50.2 kg/d) than the multiparous groups on the NL treatments (46.6 kg/d). Exposure to increased lighting enhanced milk production in multiparous cows but had no effect on primiparous animals. Feed intakes were similar for the multiparous animal groups exposed to increased day length. Since no physiological measurements were obtained, the causes of the increase in milk yield are not known, but increased lighting is an effective management practice that can be used to improve milk production on high-producing dairy herds in the San Joaquin Valley of California.

(Key Words: photoperiod, dairy cattle, supplemental light)

INTRODUCTION

Long-day photoperiod (**LDPP**, 16 to 18 h of light) is galactopoietic in dairy cows and has been shown to increase milk yield by 6.5% in dairy cows (Dahl et al., 1997; 2000). Studies involving lactating dairy cows exposed to extended light have shown that manipulation of the daily light-dark cycle increased milk yield by 2 to 3 kg/d, with little or no change in milk composition. A daily lighting regimen of 16 h of light, 8 h of darkness (16L:8D) increased milk yield 8 to 10% in Holstein cattle compared with production of cows exposed to natural Michigan winter photoperiods of 9 to 12 h of light per day (Peters et al., 1978, 1981). Photoperiod has been used successfully by a number of producers. However, the endocrine mechanism underlying this response is not known.

There does not appear to be a stage of lactation effect in response to LDPP and there is no evidence that LDPP alters mammary growth during lactation or pregnancy. The increase in milk yield due to LDPP has been shown to be additive to increased milk yields due to exogenous bovine somatotropin (**bST**) and increased milking frequency (Miller et al., 2000). Furthermore, studies evaluating effects of LDPP during the dry period on milk yield in subsequent lactation do not show any positive effects of LDPP. On the contrary, Komaragiri et al. (1997) and Aharoni et al. (2000) reported that short-day photoperiod treatment of pregnant cows resulted in greater milk yield in the subsequent lactation thus indicating that manipulation of light exposure has potential for increasing milk yield in cattle. The objective of our study was to determine if LDPP was an effective management practice that could be used to increase milk production on a high-producing dairy herd in the San Joaquin Valley of California.

MATERIALS AND METHODS

Ninety-eight multiparous and 60 primiparous Holstein animals were utilized in an extended lighting trial to investigate photoperiod's effect on milk yield. After parturition all animals were housed in one pen until 20 DIM under normal daylight/nightlight conditions. On d 21, animals were randomly assigned to two treatment groups receiving SL or NL. Diets for both treatments were balanced for 52 kg milk/d and cows were fed a total mixed ration (**TMR**) to provide 100% of Nutrient Requirements of Dairy Cattle (**NRC** 2001) requirements daily at 0600 and 1600 (Table 1).

There were two groups of 49 multiparous and two groups of 30 primiparous animals assigned to SL or NL. While in freestalls both multiparous and primiparous cows in the SL groups were exposed to 17 h of natural light and SL above 15 FC and 7 h of light below 5 FC in the freestall area. The light exposure for the NL groups followed the normal sunrise-sunset pattern common for the north 40th parallel of sunrise 0530 to 0730 and sunset 1700 to 1900, an average of 12 h of light and darkness. Light intensity was measured every 2 wk at 2200 to 2300 at two points in the freestall barn (feed manger and outside lane at animal head level) and in the milking parlor holding pen (front, middle, and back). The freestall barns were 31.7 m wide and 274 m long, providing 13.7 sq m/cow.

Measurements. Daily milk weights were measured electronically by DeLaval computer software (DeLaval, Inc., Madison, WI) for each cow's milking throughout the 305-d lactation. Monthly milk composition analysis was by Dairy Herd Improvement Association (DHIA, Provo, UT.) Pen DMI was monitored and recorded using EZFEEDTM Feed Management Software (Valley Agricultural Software, Tulare, CA).

TMR samples were obtained and wet chemistry analyses were conducted by the DHI New York Laboratory, Ithaca, NY.

Statistical Analysis. Data were analyzed using PROC MIXED procedures of SAS[®] (SAS Institute, Inc., Cary, NC). Previous 305-d mature equivalent milk yields were included as a covariate in the analysis for multiparous cows. Daily milk yields were collapsed into weekly averages. Dependent variables tested were individual cow milk yield and pen DMI. The independent variables for milk yield and DMI included treatment, time, and the respective interactions. Time (week of study) was fit as a repeated measure and treatment means were separated using the probability of a statistical difference (PDIF) option for all.

RESULTS AND DISCUSSION

There was no milk yield difference ($P = 0.81$) between primiparous animals assigned to SL (33.0 kg/d) or NL (32.9 kg/d) treatments (Figure 1). Multiparous cows in SL groups produced more ($P < 0.01$) milk (50.2 kg/d) than the multiparous groups on the NL treatments 46.6 kg/d; (Figure 2). Average pen DMI for multiparous (29.3 vs. 28.4 kg/d) and primiparous (23.4 vs. 22.4 kg/d) animals exposed to SL or NL were not measurably different. The 3.6-kg production response observed in multiparous cows is consistent with other researchers (Peters et al., 1978; 1981; Dahl et al., 2000) who reported greater milk yields from cows exposed to increased lighting regimens. The increase observed in the current study is most likely not attributed to the slight increase in DMI. However, since milk was measured on individual cows while DMI was measured on a pen basis, the effects of DMI are not possible to elucidate from the current study.

IMPLICATIONS

Exposure to increased lighting enhanced milk production in multiparous cows but had no effect on primiparous animals. Intakes were similar for the multiparous animal groups exposed to increased day length; however, milk yield was improved. Since no physiological measurements were obtained it is not known what contributed to the increase in milk yield, but increased lighting is an effective management practice that can be used to improve milk production on high-producing dairy herds in the San Joaquin Valley of California.

ACKNOWLEDGEMENTS

We would like to thank J. Fiscalini for allowing us to conduct this study and R. Mattingly for his help with the trial.

LITERATURE CITED

- Aharoni, Y., A. Brosh, and E. Ezra. 2000. Short communication: Prepartum photoperiod effect on milk yield and composition in dairy cows. *J. Dairy Sci.* 83:2779.
- Dahl, G. E., T. H. Elasser, A. V. Capuco, R. A. Erdman, and R. R. Peters. 1997. Effects of a long daily photoperiod on milk yield and circulating concentrations of insulin-like growth factor-I. *J. Dairy Sci.* 80:2784.
- Dahl, G. E., B. A. Buchanan, and H. A. Tucker. 2000. Photoperiodic effects on dairy cattle: A review. *J. Dairy Sci.* 83:885.
- Komaragiri, M. V. S., and R. A. Erdman. 1997. Factors effecting body tissue mobilization in early lactation dairy cows. 1. Effects of dietary protein mobilization of body fat and protein. *J. Dairy Sci.* 80:929.
- Miller, A. R. E., E. P. Stanisiewski, R. A. Erdman, L. W. Douglas, and G. E. Dahl. 1999. Effects of long daily photoperiod and bovine somatotropin (Trobest[®]) on milk yield in cows. *J. Dairy Sci.* 82:1716.
- NRC. 2001. Nutrient requirements of dairy cattle. 7th rev. ed. Natl. Acad. Sci., Washington, DC.
- Peters, R. R., L. T. Chapin, R. S. Emery, and H. A. Tucker. 1981. Milk yield, feed intake, prolactin, growth hormone and glucocorticoid response of cows to supplemental light. *J. Dairy Sci.* 64:1671.
- Peters, R. R., L. T. Chapin, K. B. Leining, and H. A. Tucker. 1978. Supplemental lighting stimulates growth and lactation in cattle. *Science.* 199:911.


Table 1. Ingredient and nutrient composition of experimental diets (% DM).

Item	Mature Cows	First Lactation
Ingredient		
Corn silage	22.7	21.1
Flaked corn	12.2	11.5
Flaked barley	5.2	4.9
Canola meal	13.5	14.1
Whole cottonseed	9.7	10.1
Alfalfa hay	8.9	8.7
Soy hulls	6.8	7.1
Almond hulls	6.1	6.3
Wheat millrun	3.32	3.5
Beef tallow	1.0	1.0
Sudan grass	4.3	4.5
Molasses (21%cp)	1.4	1.5
Trace min/vit premix ^a	2.5	2.6
Blood meal	2.0	2.1
Megalac ^b	0.7	1.0
Nutrient		
DM %	60.9	62.3
CP %	17.5	17.8
Rumen Undegradable Protein, % of CP	35.5	35.5
Net Energy of Lactation, Mcal/kg	1.7	1.7
Fat, %	5.8	6.2
NDF,%	36.9	36.9
ADF,%	24.4	24.5
Non-Structural Carbohydrates	33.2	32.7

^a Composition: not less than 33.3CaCO₃, 17.9 NaHCO₃, 8.1KCO₃, 6.7NaCl, 5.6MgO, 0.12Fe, 0.04Cu, 0.25Zn, .004I, 0.006Co, .001Se; contributed 7.0 KIU vit A, 1.1KIU vit D, 18.7IU vit E per kg diet Dm. Ration provided by M. A. Etchebarne, Ph.D.

^bArm & Hammer Animal Nutrition Group, Princeton, NJ.

Figure 1. Effect of photoperiod on milk yield of primiparous dairy cows



● Extended Light

○ No Light

Figure 2. Effect of photoperiod on milk yield of multiparous dairy cows

